

DECLARATION

I, Yoshiaki TODAKA of The Patent Corporate Body ARUGA PATENT OFFICE, 3-6, Nihonbashiningyocho 1-chome, Chuo-ku, Tokyo 103-0013, Japan do solemnly and sincerely declare that I well understand both Japanese and English languages, and that the attached English version is a true and complete translation of Japanese Patent Application No. 2003-197096 filed July 15, 2003 in the name of KAO CORPORATION.

Dated June 4, 2008

  
\_\_\_\_\_  
Yoshiaki TODAKA

[Name of Document] Description

[Title of the Invention] ORAL CAVITY COMPOSITION

[Claims]

[Claim 1]

An oral cavity composition comprising a first composition (A) and a second composition (B) given below, the first composition (A) and the second composition (B) being discretely packed from each other so that the two compositions can be alternately used and then come to be mixed with each other at a tooth region when applied thereto:

(A) a first composition containing a fluoride ion-supplying compound and an inorganic phosphoric acid or a salt thereof; and

(B) a second composition containing a calcium salt of organic acid, wherein an organic acid constituting the calcium salt of organic acid has a pKa value ranging from 3 to 11, or at least one pKa value ranging from 3 to 11 when the organic acid has plural pKa values.

[Claim 2]

The oral cavity composition according to claim 1, wherein an aqueous solution of the first composition (A) has a pH value ranging from 2 to 6 and/or an aqueous solution of the second composition (B) has a pH value ranging from 6 to 12.

[Claim 3]

The oral cavity composition according to the preceding claim 1 or 2, wherein said inorganic phosphoric acid is an

orthophosphoric acid.

[Claim 4]

The oral cavity composition according to any one of the preceding claims 1 through 3, wherein said calcium salt of organic acid is a calcium salt of polyol phosphate.

[Claim 5]

The oral cavity composition according to any one of the preceding claims 1 through 4, wherein said first composition (A) and/or said second composition (B) further contains a monofluorophosphate.

[Claim 6]

The oral cavity composition according to any one of the preceding claims 1 through 5, wherein the molar ratio of said fluoride ion-supplying compound to said inorganic phosphoric acid or salt thereof (fluoride ion-supplying compound/inorganic phosphoric acid or salt thereof) falls in the range of 0.1 to 10.

[Claim 7]

The oral cavity composition according to any one of the preceding claims 1 through 6, wherein said second composition (B) is a powder.

[Claim 8]

A product for dental treatment comprising a first composition (A) and a second composition (B) given below, wherein the first composition (A) and/or the second composition (B) is supported on a carrier selected from the group consisting of

paper, cloth, nonwoven fabric, absorbent cotton, sponge and porous film, and wherein the first composition (A) and the second composition (B) are discrete from each other so that the two compositions can be alternately used and then come to be mixed with each other at each tooth region when applied thereto:

(A) a first composition containing a fluoride ion-supplying compound and an inorganic phosphoric acid or a salt thereof; and

(B) a second composition containing a calcium salt of organic acid, wherein an organic acid constituting said calcium salt of organic acid has a pKa value ranging from 3 to 11, or at least one pKa value ranging from 3 to 11 when the organic acid has plural pKa values.

[Claim 9]

A method of treating teeth, comprising alternately applying a first composition (A) and a second composition (B) to a tooth:

(A) a first composition containing a fluoride ion-supplying compound and an inorganic phosphoric acid or a salt thereof; and

(B) a second composition containing a calcium salt of organic acid, wherein an organic acid constituting the calcium salt of organic acid has a pKa value ranging from 3 to 11, or at least one pKa value ranging from 3 to 11 when the organic acid has plural pKa values.

[Claim 10]

The treating method according to the preceding claim 9, wherein the first composition (A) and/or the second composition (B) is supported on a carrier selected from the group consisting of paper, cloth, nonwoven fabric, absorbent cotton, sponge and porous film.

[Claim 11]

The treating method according to the preceding claim 10, wherein the carrier supporting the first composition (A) and/or the second composition (B) is impregnated with water immediately prior to application thereof to a tooth.

[Detailed Description of the Invention]

[0001]

[Technical Field to which the Invention pertains]

The present invention relates to an oral cavity composition capable of shortening the required treatment period, efficiently preventing tooth decay, and efficiently preventing or treating dentin hyperesthesia, by applying a first composition (A) and a second composition (B) respectively having different components, alternately to a tooth.

[0002]

[Conventional Art]

As methods effective for the prevention of tooth decay, there are several means known to date, such as (1) strengthening the acid resistance of teeth to render their constituents harder to elute in acid, which represents one of the direct causes of dental caries, and (2) facilitating the remineralization process on teeth to compensate for their eluted constituents at a higher rate.

[0003]

Since fluoride acts to turn hydroxyapatite which is a major constituent of teeth into a fluoroapatite to render the constituent hard to elute in acids and is highly effective for facilitating the formation of hydroxyapatite to thereby facilitate the remineralization of teeth, it has become common at dental clinicals to coat teeth with a fluoride as treatment for preventing dental caries. In such cases, the most commonly

used fluoride formulation is an acidulated phosphate fluoride solution (hereinafter referred to as APF).

[0004]

However, the APF treatment takes a long time to complete the formation of calcium fluoride, because it includes a process which facilitates the double decomposition of teeth by the action of an acid and then exploits calcium contained there to produce calcium fluoride for fluoride uptake into the teeth. Specifically, for this method of treatment, it has been necessary that APF is first applied onto human teeth to be left as it is for 4 minutes, and further, the eating and/or drinking is forbidden for additional 30 minutes or longer. In this case, a person undergoing the care or treatment is required to exercise patience in keeping his or her mouth open for 4 minutes after APF application, and further, such a patient subjected to the treatment may feel rather burdensome because APF has a nasty taste due to its strongly acidic taste. Especially, for patients who are children, such a bad taste might be more burdensome.

[0005]

On the other hand, as periodontal disease patients increase in number, dentin hyperesthesia caused by a stimulus occurring when eating and/or drinking something hot or cold and transferred directly to pulp nerves due to exposed dentin at dental roots has now become a problem to be seriously dealt with. To prevent or treat this dentin hyperesthesia, it is necessary to block the transmission of external stimulus by heat, acids, or etc.

to pulp nerves by obturating dental canaliculi present in dentin.

[0006]

While it has so far been done as preventive and curative treatment for dentin hyperesthesia to apply a 5% sodium fluoride varnish coating onto teeth, such application treatment is not free from problems yet, in that this method uses calcium contained in teeth and saliva to obturate the dental canaliculi and it takes time for calcium to turn to its fluoride and thus a longer time elapses before the dental canaliculi is effectively obturated, or in that the effect of the treatment has a low persistence after application (cf. the nonpatent literature 1.)

[0007]

The patent literature 1 given below discloses a product for the remineralization of dental enamel, containing a first separate ingredient containing a water-soluble calcium salt, and a second separate ingredient containing a phosphate and at least one water-soluble fluoride salt that produces fluorine ions. However, the mineralization product disclosed in the patent literature 1 is taught to be "applied as a mixture prepared by mixing the first and second solutions together" (cf. the patent literature 1, page 19, lines 3 to 4).

[0008]

In this regard, in a case where the first and the second liquids are first mixed together and then the resultant mixture is applied on to the teeth, there has been a problem that it is difficult to form calcium phosphates selectively on a tooth



surface, because if the first solution containing a water-soluble calcium salt is mixed with the second solution comprising a phosphate and a water-soluble fluoride salt, such calcium phosphates (including partially fluorinated salts) are formed in the agent during the course of mixing.

[0009]

[Patent document 1]

(JP-A-2001-523217).

Nonpatent literature 1: "Tooth Wear and Dentin Hyperesthesia" translated by Kenichi Kobayashi, et al., p.357, Ishiyaku Publishers, Inc., 2003

[0010]

[Problems to be solved by the Invention]

The present invention has been made to overcome the aforementioned drawbacks of the prior art, and it is an object of the invention to provide an oral cavity composition exhibiting a good effect for dental cavity prevention and excellent preventive and therapeutic effects for dental hyperesthesia, even when used in a treatment taking only some ten seconds, and a method of treatment using such a composition.

[0011]

[Means for solving the Problems]

The present invention provides an oral cavity composition containing a first composition (A) and a second composition (B) given below, said first composition (A) and said second composition (B) being discretely packed from each other so that

said two compositions can be alternately used and then come to be mixed with each other at each tooth region when applied thereto:

(A) a first composition containing a fluoride ion supplying compound and an inorganic phosphoric acid or a salt thereof; and

(B) a second composition containing a calcium salt of organic acid, wherein an organic acid constituting said calcium salt of organic acid has a pKa value ranging from 3 to 11, or at least one pKa value ranging from 3 to 11 when said organic acid has plural pKa values.

[0012]

The application of the oral cavity composition according to the present invention permits an efficient uptake of a quantity of a fluoride into teeth with a brief treatment, while providing a high preventive effect against dentin hyperesthesia by calcium fluoride and calcium phosphate particles that are produced in the course of treatment and act to obturate the dental canaliculi. Further, of the oral cavity composition according to the present invention, the second composition (B), when applied to a tooth after application of the composition (A), exerts a buffer action to increase the pH value of the first composition (A) so as to reduce the acidic taste caused by the first composition (A), thus leading to an improved feeling of use as an oral cavity composition.

[0013]

Besides, the present invention has permitted an efficient

uptake of a fluoride in a shorter time period by using the first composition (A) and the second composition (B) as aqueous solutions thereof, with the former aqueous solution having a pH value ranging from 2 to 6 and/or the latter aqueous solution having a pH value ranging from 6 to 12. Furthermore, it may be possible to form calcium phosphate from phosphate ions contained in the first composition (A) or the tooth and calcium derived from the second composition (B) and then fix the resultant calcium phosphate onto the tooth surface. At the same time, calcium fluoride may be fixed on the tooth surface.

[0014]

With the pH value of the first composition (A) below 6, a double decomposition reaction forming calcium fluoride from hydroxyapatite and a fluoride is facilitated and as a result the fluorine uptake into the tooth increases to cause the fluoride to be taken into the tooth as calcium fluoride. While, if the pH value is below 2, the calcium fluoride gets dissolved, resulting in a reduced fluorine uptake. Therefore, it is preferred that the first composition (A) according to the present invention has a pH value in the range of 2 to 6 in its aqueous solution. More preferably, the pH value ranges from 3 to 5.5 and further preferably from 4 to 5.

[0015]

On the contrary, since the second composition (B) is used for neutralizing the first composition (A) to facilitate the aggregation and fixation of the calcium fluoride or calcium

phosphate, it is preferred that the second composition (B) has a pH value ranging from the neutral to the alkali zone in an aqueous solution thereof. Specifically, the pH value of the second composition (B) ranges from 6 to 12, more preferably from 7 to 11 and further preferably from 7 to 9.

[0016]

The first composition (A) according to the present invention is applied onto a tooth to permit the supply of fluorine ions and phosphoric acid in order to form calcium fluoride and calcium phosphate on the tooth surface.

[0017]

The second composition (B) according to the present invention is used to externally supply calcium for permitting the quick formation of calcium fluoride and consequently enabling the teeth treatment to be accomplished in a shorter time period (some ten seconds), thus substantially relieving the burden on persons (especially children) under dental treatment. Besides, the concurrent formation of calcium fluoride with calcium phosphate improves the acid resistance of teeth and facilitates the remineralization of teeth, and also highly improves the effect to obturate the dental canaliculi. Further, since the second composition (B) according to the present invention has a buffering action, it is possible to increase the pH value of the first composition (A) to reduce the acidic taste caused by the first composition (A) and thus improve the feeling of use of the oral cavity composition.

[0018]

Further, according to the present invention, the first composition (A) and the second composition (B) are alternately applied directly onto a tooth so that the tooth surface serves as a site where the first composition (A) and the second composition (B) are reacted with each other, which allows the aggregation of calcium fluoride or calcium phosphate to occur limitedly on the tooth surface, and consequently the present invention improves the effect for improving the uptake of calcium fluoride and calcium phosphate into teeth and thus provides a high effectiveness in dental cavity prevention and also excellent preventive and treatment effects against dental hyperesthesia.

[0019]

Moreover, APF itself has not been recognized to have any preventive effect against dentin hyperesthesia. When the first composition (A) containing APF and the second composition (B) according to the present invention are alternately applied to a tooth, APF cooperates with the second composition (B) containing a calcium salt of an organic acid to efficiently form calcium fluoride or calcium phosphate on the surface of the tooth, thus making it possible to obtain not only a high preventive effect against dental cavity but also excellent preventive effects against dentin hyperesthesia and the like.

[0020]

Now, the efficient formation of calcium fluoride and calcium phosphate on the tooth surface realized by the

cooperation of the first composition (A) and the second composition (B) according to the present invention will be described more specifically. After supplying a sufficient quantity of the first composition (A) to a target tooth region under treatment, the second composition (B) is supplied by application to the target tooth region in an excess quantity sufficient for the entirety of the first composition (A) to complete its reaction with the second composition (B). Then, the first composition (A) is supplied again to the target teeth region in an excess quantity sufficient for the entirety of the residual second composition (B) to complete the first composition (A). By repeating such a cycle, the present invention permits the first composition (A) and the second composition (B) to cooperate with each other to efficiently form calcium fluoride or calcium phosphate on the tooth surface.

[0021]

[Mode for Carrying out the Invention]

The fluoride ion-supplying compound according to the present invention is not specifically limited as long as it is a material usable in the oral cavity, and examples thereof include inorganic fluorides such as sodium fluoride, potassium fluoride, ammonium fluoride, lithium fluoride, etc., and organic fluorides such as amine fluorides, etc. Among them, sodium fluoride, sodium monofluorophosphate, tin fluoride, lithium fluoride and ammonium fluoride are preferred in respect of safety, solubility, flavor, etc..

[0022]

It is to be noted here that monofluorophosphates (e.g., sodium monofluorophosphate, potassium monofluorophosphate, ammonium monofluorophosphate, or etc.) may be used in combination with the above-described fluoride ion-supplying compound and calcium salt of organic acid.

If the calcium fluoride and calcium phosphate produced from the composition (A) and the composition (B) according to the present invention contain a monofluorophosphate such as sodium monofluorophosphate, the acid resistance of the teeth will be further improved. The first composition (A) and/the second composition (B) according to the present invention may contain sodium monofluorophosphate in a quantity ranging from 0.01 to 20 wt%, preferably from 0.05 to 5 wt% and more preferably from 0.05 to 2 wt%.

[0023]

It is preferred that the first composition (A) of the present invention has a fluorine ion concentration in the range of 50 ppm to 200 ppm or 0.0025 mol to 1 mol. More preferably, the fluorine ion concentration ranges from 500 ppm to 10000 ppm or 0.025 mol to 0.5 mol and particularly preferably from 1000 ppm to 9000 ppm or 0.05 mol to 0.5 mol.

[0024]

According to the present invention, the inorganic phosphoric acids or salts thereof include, for example, alkali metal salts and ammonium salts of orthophosphoric acids such

as potassium orthophosphate, sodium orthophosphate, or ammonium orthophosphate, and etc., monopotassium phosphate, dipotassium phosphate, tripotassium phosphate, monosodium phosphate, disodium phosphate and trisodium phosphate, and pyrophosphoric acids and salts thereof (e.g., sodium pyrophosphates, potassium pyrophosphates, ammonium pyrophosphates, or etc.). The inorganic phosphoric acid concentration of the second composition (B) of the present invention ranges preferably from 0.005 mol to 3 mol, more preferably from 0.05 mol to 1 mol and particularly preferably from 0.1 mol to 0.5 mol.

[0025]

The calcium salt of an organic acid according to the present invention has a pKa value of 3 to 11 if the salt is a compound having a single pKa value, or is a calcium salt of an organic acid having at least one pKa value of 3 to 11 if the salt has a plurality of pKa values. Specifically, such calcium salts of an organic acid include calcium salts of polyol phosphates such as glycerophosphate, glyceryl aldehyde-3-phosphate, erythrose-4-phosphate, ribose-5-phosphate, glucose-1-phosphate, glucose-6-phosphate, inositol monophosphate, inositol hexaphosphate, fructose-1-phosphate, fructose-6-phosphate, fructose-1,6-diphosphate, ascorbate-2-phosphate, phosphorylated maltotriose, phosphorylated maltotetraose, and etc.

[0026]

Also, the calcium salt of organic acids according to the



present invention include water-soluble compounds composed of organic acids and calcium such as calcium lactate, calcium acetate, calcium gluconate, calcium benzoate, calcium glycerophosphate, calcium formate, calcium fumarate, calcium butyrate, calcium isobutyrate, calcium malate, calcium maleate, calcium propionate, and etc.

[0027]

It is preferred that the first composition (A) according to the present invention be used in the form of a liquid, paste or gel. It is also preferred that the molar ratio of the fluoride ion-supplying compound vs. the inorganic phosphoric acid or salt thereof (fluoride ion-supplying compound/inorganic phosphoric acid or salt thereof) of the first composition (A) according to the present invention falls in the range of 0.1 to 10.

[0028]

According to the present invention, the second composition (B) is not limited to any particular state but may take any state that can supply calcium ions and permit it to mix quickly with the first composition (A) and, preferably, it is provided in the form of a liquid, paste, gel or powder. The content of the calcium salt of organic acid of the second composition (B) ranges preferably from 0.005 mol to 1 mol, more preferably from 0.01 mol to 0.5 mol and particularly preferably from 0.02 mol to 0.2 mol.

[0029]

The second composition (B) according to the present

invention may be used in the state of a powder or it may be supported in a state substantially free of water content on a suitable carrier. For example, the second composition (B) according to the present invention may be supported (by adhesion or adsorption) on a carrier and water may be added to the carrier in use, resulting in that a high calcium concentration and buffer action can be maintained at the site of treatment action (e.g., tooth surface subjected to treatment) and the ease of use can be further improved.

[0030]

In cases where the first composition (A) or the second composition (B) is a powder, the powder content may range preferably from 0.1 to 100 times, more preferably from 0.2 to 50 times and particularly preferably from 0.2 to 20 times the mass of the carrier used.

The first composition (A) and the second composition (B) according to the present invention both may be in the form of liniment, mouth wash or dentifrice, or either one may be in the form of mouth wash and the other dentifrice, or vice versa.

[0031]

According to the present invention, the molar ratio of the fluoride ion-supplying compound of the first composition (A) vs. the calcium salt of organic acid of the second composition (B) is not specifically limited, but the molar ratio may range preferably from 0.02 to 50, more preferably from 0.05 to 20 and particularly from 0.2 to 5.

[0032]

The carrier for the first composition (A) and/or the second composition (B) according to the present invention is not specifically limited, but may be any appropriate water absorbent material having a sufficient strength for application to the tooth surface. More specifically, appropriate materials for the carrier include paper, cloth (e.g., gauze, or etc.), nonwoven fabrics (e.g., of ultrafine fibers, natural fibers, or etc.), absorbent cotton, sponges, porous films, and etc.

The above-described porous films may include celluloses, acetates, polyurethanes, porous synthetic rubbers, collagens, and etc.

[0033]

Specifically, to treat a tooth by alternately applying thereto the first composition (A) and the second composition (B) according to the present invention in cases where the compositions are provided as a liquid liniment, the first composition (A) may be applied to the target site and left to stand for 5 to 10 seconds thereafter and then the second composition (B) may be applied to the target site and left to stand for 5 to 10 seconds thereafter, as well. Such a cycle may be repeated one or more times. Alternatively, the above-described process may be accomplished in reverse order, that is, the second composition (B) may first be applied to the target site and left to stand for 5 to 10 seconds thereafter and then the first composition (A) is applied to the target site

to be left to stand for 5 to 10 seconds thereafter.

[0034]

Also, where the first composition (A) is a mouth wash and the second composition (B) is a dentifrice, the mouth may be rinsed for 30 seconds with the mouth wash of the first composition (A) and then the teeth may be brushed for 3 minutes with the dentifrice of the second composition (B), and such a cycle may be repeated one or more times. Further, the teeth may first be brushed with the dentifrice of the second composition (B) for 3 minutes, followed by 30 seconds of mouth rinsing with the mouth wash of the first composition (A).

[0035]

In cases where the first composition (A) and the second composition (B) are both provided as mouth washes, the mouth may first be rinsed for 30 seconds with the mouth wash of the first composition (A) and then rinsed for another 30 seconds with the mouth wash of the second composition (B), and such a cycle may be repeated one or more times. In use, the second composition (B) may be applied first, and then the first composition (A). When the first composition (A) is provided in the form of liniment and the second composition (B) in the form of powder, the liniment of the first composition (A) may first be applied to the target site and left to stand for 5 to 10 seconds and then the powder of the second composition (B) may be applied to the target site to be left to stand for 5 to 10 seconds thereafter. Such a cycle may be repeated one or more

times. In use, the second composition (B) may be applied first, and then the first composition (A), as well.

[0036]

In addition to the aforementioned ingredients, the oral cavity composition of the present invention may contain, for example, a foaming agent, foaming aid, abrasive, wetting agent, binder, extender, sweetener, preservative, bactericide, medicinal ingredient, self-adhesive, pigment, colorant, flavor, or etc., as appropriate. Besides, there is no limitation to the use of polyethylene glycol or like skin-lightening ingredients in combination with the oral cavity composition according to the present invention.

Furthermore, the oral cavity composition of the present invention may be prepared in the dosage form of a solution, gel, paste, or etc., for example, and may be used as a dentifrice, liquid dentifrice, mouth wash, or etc.

[0037]

[Example]

The percentage in the Examples indicates percentage by weight.

Example 1     Dental treating agent 1

A	Sodium fluoride	2%
	Phosphoric acid	1.6%
	Purified water	balance
	pH	3.6
B	Calcium glycerophosphate	3%

Sodium monofluorophosphate	1%
Purified water	balance

Usage: Composition A is applied first and then the composition B.

[0038]

#### Example 2 Dental treating agent 2

A	Sodium fluoride	2%
	Phosphoric acid	3%
	Purified water	balance
	pH	adjusted to pH 4.0 with sodium hydroxide
B	Sodium glycerophosphate	50%
	Xylitol	20%
	Glycerin	10%
	Sodium monofluorophosphate	1%
	Dextrin	balance

The ingredients are mixed together so that the total weight amounts to 50% of the swab weight. After applying the composition A, the composition B is wetted with water and applied

[0039]

#### Example 3 Dental treating agent 3 (applied and rinsed away from the mouth)

A	Sodium fluoride	2%
	Phosphoric acid	2%
	Sodium monofluorophosphate	1%
	Purified water	balance
	Adjusted to pH 3.5 with sodium hydroxide	

B	Calcium lactate	1%
	Xylitol	10%
	Polyoxyethylene hydrogenated castor oil	1%
	Flavor	0.1%
	Purified water	balance

After applying the composition A, the composition B is used to rinse the mouth.

[0040]

#### Example 4 Dentifrice and mouth wash

A	Sodium fluoride	0.21
	Phosphoric acid	0.5
	Xylitol	10
	Sorbitol solution (70%)	40
	Polyethylene glycol 600	5
	Saccharin	0.1
	Sodium lauryl sulfate	1
	Silicic acid anhydride	15
	Carboxymethylcellulose sodium salt	1
	Flavor	1
	Purified water	balance

Adjusted to pH 5 with sodium hydroxide.

B	Calcium glycerophosphate	1%
	Sodium monofluorophosphate	0.14%
	Xylitol	10%
	Polyoxyethylene hydrogenated castor oil	1%
	Flavor	0.1%

Purified water balance

After brushing teeth with the dentifrice of the composition A, the mouth is rinsed with the mouth wash of the composition B.

[0041]

Example 5 Two mouth washes

A	Sodium fluoride	0.05
	Phosphoric acid	0.2
	Sorbitol solution (70%)	10
	Sodium lauryl sulfate	0.2
	Flavor	0.1
	Purified water	balance

Adjusted to pH 4.5 with sodium hydroxide.

B	Calcium glycerophosphate	1%
	Sodium monofluorophosphate	0.14%
	Xylitol	10%
	Polyoxyethylene hydrogenated castor oil	1%
	Flavor	0.1%
	Purified water	balance

Usage: The mouth is rinsed with the mouth wash of the composition A, followed by rinsing with the B mouth wash.

[0042]

<Comparative examples>

Comparative example 1 Dental liniment (acidulated phosphate fluoride solution APF)

Ingredients	Sodium fluoride	2%
	Phosphoric acid	1.5%



Comparative example 2     Dentin desensitizer (F Varnish, Bee Brand Medico Dental. Co., Ltd.)

Ingredient	Sodium fluoride	5%
------------	-----------------	----

Comparative example 3     Dentin hyperesthesia preventive dentifrice (Shumitect, Earth Chemical Co. Ltd.)

Active ingredient	Potassium nitrate
-------------------	-------------------

[0043]

Comparative example 4

A	Sodium fluoride	0.21
	Xylitol	10
	Sorbitol solution (70%)	40
	Polyethylene glycol 600	5
	Saccharin	0.1
	Sodium lauryl sulfate	1
	Silicic acid anhydride	15
	Carboxymethylcellulose sodium salt	1
	Flavor	1
	Purified water	balance

[0044]

<Evaluation of fluorine uptake>

1. Determination of fluorine adsorption on HAP pellets

In the Examples 1 through 5, HAP pellets (10 mm × 10 mm × 2 mm) each were treated for 10 seconds with 10 ml of A, and then with 10 ml of B for another 10 seconds. This process was repeated three times with the total treating time amounting to 1 minute for each HAP pellet. For the comparative example 1, the HAP

pellet was immersed in its 10 m% solution for 4 minutes. The composition of the comparative example 2 was applied to the HAP pellet and retained in the applied state for 1 hour before removal. For the comparative examples 3 and 4, the pellets each were immersed in the respective dentifrices for 3 minutes. The fluoride adsorbed on the surface of each HAP pellet was extracted with hydrochloric acid, and using an ion analyzer (Expandable ionAnalyzer EA 940 (made by ORION company)) with fluoride ion electrodes (ionplus-Fluoride (made by ORION company)) fluorine adsorbed on the surface of HAP pellet was determined.

[0045]

## 2. Results

The results of the evaluation are shown in Table 1. The Examples 1 through 3 achieved higher fluorine uptakes to teeth in a treating time as short as 1 minute. On the contrary, the comparative example 1 (representing the existing level of the art) took as long as 4 minutes to get the same level of fluorine uptake as achieved by the preferred examples 1 through 3, thus demonstrating the effectiveness of the present invention for the fluorine uptake in a shorter time period. The preferred examples 4 and 5 achieved the fluorine uptakes of 0.6 and 0.3  $\mu\text{g}/\text{cm}^2$ , respectively, in a 1 minute treatment, although the comparative example 4 representing common dentifrices got only a small fluorine uptake of 0.1  $\mu\text{g}/\text{cm}^2$  in spite of its treating time as long as 3 minutes, that is, the preferred examples 4 and 5 of the present invention showed fluorine uptakes several

times greater than the comparative example 4.

[0046]

	Treating time	Fluorine uptake ( $\mu\text{g}/\text{cm}^2$ )
Example 1	1 min.	5.2
Example 2	1 min.	5.8
Example 3	1 min.	4.9
Example 4	1 min.	0.6
Example 5	1 min.	0.3
Comparative example 1	4 min.	4.2
Comparative example 2	1 hr.	2.2
Comparative example 3	3 min.	Below detectable level
Comparative example 4	3 min.	0.1

[0047]

<Evaluation of efficacy on dentin hyperesthesia>

#### 1. Methodology

One hundred persons aged from 24 to 64 years suffering from dentin hyperesthesia were employed as subjects for evaluation. The subjects were divided into five groups of 20 subjects each and the subjects belonging to three groups thereof were treated for 10 seconds with the solutions of the preferred examples 1 through 3 in 3 cycle alternate treatment mode, respectively, while for the comparative examples the subjects of the remaining one group were treated for 4 minutes with APF commonly used for dental cavity prevention (comparative example 1) and the subjects of another remaining one group were treated by applying to them a 5% sodium fluoride preparation widely used in the past for

the treatment of dentin hyperesthesia (comparative example 2). The efficacy of each example for dentin hyperesthesia was evaluated twice, namely, immediately after the application of the medicament of each example and 1 week thereafter, based on any changes in rational symptoms observed in each subject.

[0048]

## 2. Results

The results of evaluation immediately after the alternate treatment with the first composition (A) and the second composition (B) of the present invention are shown in Table 2 along with the results obtained from the corresponding evaluation of the comparative examples. Most subjects having been suffering from dental hyperesthesia showed the disappearance of such symptoms of dental hyperesthesia after undergoing the respective alternate treatment for 1 minute with Examples 1 through 3 and they became free from hypersensitive toothache (Example 1: for 19 subjects among 20). To the contrary, the comparative example 1 showed a result that 15 subjects out of 20 did not undergo any improvement in their dental hyperesthesia symptoms even after 4 minutes of treatment and their hypersensitive toothache remained as before. Besides, as shown in Table 1, although the comparative example 1 exhibited a high fluorine uptake into teeth, its effectiveness for improvement of dentin hyperesthesia was very low.

[0049]

Further, in the case of the comparative example 2

representing a formulation commonly used for the treatment of dentin hyperesthesia (nonpatent literature 1), dental hyperesthesia disappeared in 11 subjects out of 20 and an improvement of the hypersensitive toothache symptom was observed in 8 subjects, but no such improvement of dental hyperesthesia symptom was observed in 3 subjects. From this result, it was demonstrated that Examples 1 through 3 of the present invention are highly effective for improving dentin hyperesthesia.

[0050]

Further, in the comparative example 2, the formulation was required to be retained in a state as it was applied onto the tooth surface for 1 hour after the application, consequently imposing the subject under treatment to not a little burden. On the contrary, the present invention requires only as short a time as 1 minute for treatment and has demonstrated realization of a substantial reduction of the burden imposed on subjects under treatment.

[0051]

The results of the evaluation obtained one week after the alternate treatment with the first composition (A) and the second composition (B) of the present invention are shown in Table 3 along with the results obtained from the corresponding evaluation on the comparative examples. The preferred examples 1 through 3 counted 10, 12 and 12 subjects out of 20, respectively, who could experience the disappearance of dentin hyperesthesia symptom and retain their state free of the symptom thereafter,

while the comparative example 2 had only 3 subjects who could likewise experience such disappearance of dentin hyperesthesia symptom and retain their state free of the symptom thereafter. Further, the preferred examples 1 through 3 counted 4, 3 and 1 subjects out of 20, respectively, who had dental hyperesthesia symptoms 1 week after the treatment, while the comparative example 2 counted as many as 7 such subjects out of 20. From the results above, it can be said that the present invention is effective for improving the persistence of the effect of treatment against dentin hyperesthesia.

[0052]

In the alternate treatment with the first composition (A) and the second composition (B) in the preferred embodiments according to the present invention, fluoride ions and phosphoric acid were supplied by the first composition (A) and calcium ions were supplied by the second composition (B), and these ions served to form calcium fluoride and calcium phosphate at the same time on the tooth surface under treatment and in dental canaliculi so as to yield a higher treatment effect in a shorter treating time. Further, the calcium phosphate and calcium fluoride coexisting in dental canaliculi acted to promote dentin remineralization to obtrude the dental canaliculi and further provided the effect for reforming the teeth surface to improve its acid resistance and improving the persistence of the effect of treatment against dentin hyperesthesia. This demonstrated that the present invention is effective for improving dentin

hyperesthesia and for preventing its recurrence.

[0053]

On the contrary, in the comparative example 2, the agent used supplies only fluoride ions and calcium fluoride is formed using calcium ion supplied internally of the tooth, and further not only the tooth treatment takes time due to a high solubility of calcium fluoride, but also the persistence of treatment effect is inferior to the Examples.

[0054]

[Table 2]

## &lt;Results immediately after treatment&gt;

	Subjects without symptom improvement (persons)	Subjects with symptom relief (persons)	Subjects symptom disappearance (persons)
Example 1	0	1	19
Example 2	0	2	18
Example 3	0	1	19
Comparative example 1	15	5	0
Comparative example 2	3	8	11

[0055]

[Table 3]

## &lt;Results 1 week after treatment&gt;

	Subjects without symptom improvement (persons)	Subjects with symptom relief (persons)	Subjects symptom disappearance (persons)
Example 1	4	6	10
Example 2	3	5	12
Example 3	1	7	12
Comparative example 1	16	4	0
Comparative example 2	7	10	3

[0056]

Further, as a result of a questionnaire on feeling of use to the subjects who underwent the treatment, it was revealed that the Examples hardly caused any problems, but many subjects treated with the comparative example 1 answered to have felt uncomfortable due to a strong acidic taste (see Table 4).

[Table 4]



## &lt;Feeling of use&gt;

	No uncomfortabl e feeling	Somewhat uncomfortabl e feeling	Uncomfortable feeling experienced
Example 1	17	3	0
Example 2	16	4	0
Example 3	17	3	0
Comparative example 1	16	4	8
Comparative example 2	11	9	0

[0058]

<Evaluation of efficacy of dentifrices and mouth washes  
on dentin hyperesthesia>

[0059]

Eighty persons aged from 24 to 64 years suffering from dentin hyperesthesia were employed as subjects for evaluation. The subjects were divided into four groups of 20 subjects each and, in the preferred example 4, 20 subjects belonging to one group brushed their teeth with the dentifrice for 3 minutes and then rinsed the mouth with the mouth wash, twice in one day. In the preferred example 5, another 20 subjects rinsed their mouths alternately with the mouth washes A and B each three times for 10 seconds each, twice in one day. In the comparative example 3, another 20 subjects brushed their teeth with the dentifrice for 3 minutes, twice in one day. In the comparative example 4, another 20 subjects brushed their teeth with the dentifrice for 3 minutes, twice in one day. One week after, each subject was checked for any changes in rational symptoms.

[0060]

Consequently, 60% or more of subjects treated with the preferred examples experienced the disappearance of pain, as compared with 20% or less for the comparative examples, as shown in Table 5.

[0061]

<After one week>

	Subjects without symptom improvement (persons)	Subjects with symptom relief (persons)	Subjects symptom disappearance (persons)
Example 4	1	7	12
Example 5	2	4	14
Comparative example 3	10	6	4
Comparative example 4	8	9	3

[0062]

[Effect of the Invention]

The alternate application of the first composition (A) and the second composition (B) according to the present invention permits an efficient uptake of a quantity of a fluoride into teeth with brief treatment, while providing a high preventive effect against dentin hyperesthesia by dental canaliculi calcium fluoride and calcium phosphate particles that are produced in the course of treatment and act to obturate the dental canaliculi. Further, of the oral cavity composition according to the present invention, the second composition (B), when applied to teeth after application of the composition (A), exerts a buffer action to increase the pH value of the first composition (A) so as to reduce the acidic taste caused by the first composition

(A) , thus leading to an improved feeling of use as an oral cavity composition.

[Name of Document] ABSTRACT

[Summary]

[Problem] The present invention provides an oral cavity composition with which brief treatment using the same yields a high effect for dental cavity prevention and an excellent dental hyperesthesia preventive and treatment effects, and a method of treatment using such a composition.

[Solving Means] The oral cavity composition according to the present invention contains a first composition (A) containing a fluoride ion-supplying compound, an inorganic phosphoric acid and a salt thereof, and a second composition (B) containing a calcium salt of organic acid, with an organic acid constituting said calcium salt of organic acid having a pKa value ranging from 3 to 11, or at least one pKa value ranging from 3 to 11 when said organic acid has plural pKa values, wherein said first composition (A) and said second composition (B) are discretely packed from each other so that said two compositions can be alternately used and then come to be mixed with each other at a tooth region when applied thereto.

[Selected Diagram] FIG. 1